



BEEF FINISHER

MEDICATED

ACTIVE DRUG INGREDIENT

Rumensin(as Monensin Na)

22.40 gm/t

GUARANTEED ANALYSIS

Crude Protein	Min	12.00 %	Sodium (Na)	Min	0.20 %
Crude Fat	Min	5.20 %	Sodium (Na)	Max	0.27 %
Crude Fiber	Max	10.60 %	Salt (NaCl)	Min	0.10 %
ADF	Max	14.65 %	Salt (NaCl)	Max	0.14 %
Calcium (Ca)	Min	0.60 %	Potassium (K)	Min	0.80 %
Calcium (Ca)	Max	0.80 %	Vitamin A	Min	3300 IU/lb
Phosphorous (P)	Min	0.30 %			

INGREDIENTS

Steam Flaked Corn, Wheat Millrun, Cane Molasses, Dried Plain Beet Pulp Shreds, Steam Rolled Barley, Whole Cottonseed, Ground Soybean Hulls 5.26%, Soybean Meal, Safflower Meal, Corn Distillers Dried Grains With Solubles, Suncured Alfalfa Meal, Ground Corn, Hydrolized Animal and Vegetable Fat, Gelatinized Corn Flour (Corn Starch), Ground Limestone, Calcium Salts of Long-Chain Fatty Acids, Glycerin, Vegetable Oil, Sodium Sesquicarbonate, Linseed Meal, Lignin Sulfonate, Recleaned Whole Oats, Dried Whey, Sodium Sulfate, Salt, Sodium Sulfite, Rice Bran, Soybean Oil, Potassium Sulfate, Magnesium Oxide, Monocalcium Phosphate, Propionic Acid (Preservative), Brewers Dried Yeast, Dried Saccharomyces Cerevisiae Fermentation Product, Yeast Culture, Sodium Bentonite, Hydrated Sodium Calcium Aluminosilicate, Potassium Chloride, Aspergillus Oryzae Fermentation Extract, Manganese Sulfate, Natural and Artificial Flavors, Magnesium Chloride, Magnesium Sulfate, Zinc Sulfate, Iron Oxide, Propylene Glycol, Copper Sulfate, Extracted Streptomyces Meal, Ferrous Sulfate, Condensed Fermentation Corn Extractives, Vitamin A Supplement, Kelp Meal, Zinc Oxide, Manganous Oxide, Certified Color Added, Vitamin D3 Supplement, Vitamin E Supplement, Ferrous Carbonate, Manganese Proteinates, Zinc Proteinates, Copper Carbonate, Riboflavin Supplement, Copper Amino Acid Complex, Vitamin B-12 Supplement, Calcium Pantothenate, Cobalt Glucoheptonate, Biotin, Potassium Sulfite, Choline Chloride, Selenium Yeast, Corn Cob Fractions, Niacin Supplement, Mineral Oil, Calcium Iodate, Cobalt Sulfate, Pyridoxine Hydrochloride, Thiamine Mononitrate, Cobalt Carbonate, Cobalt Amino Acid Complex, Ethylenediamine Dihydriodide, Folic Acid, Sodium Selenite, Ethoxyquin (Preservative), Silicon Dioxide, Citric Acid (Preservative), Tertiary Butyl Hydroquinone (Preservative).

FEEDING DIRECTIONS

SHOW CATTLE:

FOR INCREASED RATE OF WEIGHT GAIN (Growing cattle on pasture or in dry lot [stocker and feeder, and dairy and beef replacement heifers] AND Mature reproducing beef cows [on pasture in dry lot]): Hand feed or mix in total ration so as to supply 4.48 to 17.84 lbs. of this feed per head per day, thereby supplying 50 - 200 mg monensin per head daily. During first 5 days, cattle should receive no more than 100 mg monensin per day (8.92 lbs. of this feed).

CONTINUED ON BACK --->

SHOWTIME BEEF FINISHER

(Back Side)

FEEDING DIRECTIONS (Continued)

FOR THE PREVENTION AND CONTROL OF COCCIDIOSIS CAUSED BY EIMERIA BOVIS AND E. ZUERNII (Cattle fed in confinement for slaughter): Feed, depending upon severity of challenge, 1.25 to 3.75 lbs of this feed per 100 lbs body weight daily, supplying 14 to 42 mg monensin per 100 lbs body weight daily. (Growing cattle on pasture or in dry lot [stocker and feeder, and dairy and beef replacement heifers] AND Mature reproducing beef cows [on pasture in dry lot]): Feed, depending upon severity of challenge, 1.25 to 3.75 lbs of this feed per 100 lbs body weight daily, supplying 14 to 42 mg monensin per 100 lbs body weight daily up to 200 mg monensin (17.85 lbs of this feed) per head daily. During first 5 days of feeding, cattle should receive no more than 100 mg monensin (8.92 lbs of this feed) per day.

(Calves[excluding veal calves]): Feed, depending upon severity of challenge, 1.25 to 8.92 lbs of this feed per 100 lbs body weight daily, supplying 14 to 100 mg monensin per 100 lbs body weight daily up to 200 mg monensin (17.85 lbs of this feed) per head daily.

FOR IMPROVED FEED EFFICIENCY (Cattle fed in confinement for slaughter): Feed continuously so as to supply 4.48 to 32.14 lbs. of this feed per head daily, thereby supplying 50 to 360 mg monensin per head daily.

CAUTION: Do not allow horses or other equines access to feed containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin-medicated cattle feed is safe for use in cattle only. Consumption by unapproved species may result in toxic reactions. Do not exceed the levels of Monensin recommended by the feeding directions, as reduced average daily gains may result. A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal.

REMINDER: Feed is perishable. Store this product in a cool, dry area away from insects. DO NOT feed moldy or insect-infested feed.

WARNING! DO NOT FEED TO SHEEP. This feed contains added copper at levels that may be toxic to sheep.

Net Weight: **50 Pounds (22.7 Kilograms)**

Manufactured By :

Farmers Warehouse Company
P.O. Box 160
Keyes, California 95328
(209)632-2333

15565504.02
'010-A-5819

426D