



HEIFER & STEER DEVELOPER

MEDICATED

ACTIVE DRUG INGREDIENT

Rumensin(as Monensin Na) 27.60 gm/t

Crude Protein	Min	13.00 %	Phosphorous (P)	Min	0.35 %
Crude Fat	Min	4.80 %	Sodium (Na)	Max	0.26 %
Crude Fiber	Max	13.00 %	Salt (NaCl)	Min	0.08 %
ADF	Max	18.45 %	Salt (NaCl)	Max	0.13 %
Calcium (Ca)	Min	0.50 %	Potassium (K)	Min	0.80 %
Calcium (Ca)	Max	0.70 %	Vitamin A	Min	2500 IU/lb

INGREDIENTS

Steam Flaked Corn, Whole Cottonseed, Wheat Millrun, Dried Plain Beet Pulp Shreds, Steam Rolled Barley, Cane Molasses, Safflower Meal, Soybean Hull Pellets 4.6%, Steam Rolled Oats, Corn Distillers Dried Grains With Solubles, Soybean Meal, Ground Limestone, Ground Corn, Gelatinized Corn Flour (Corn Starch), Glycerin, Vegetable Oil, Sodium Sesquicarbonate, Linseed Meal, Recleaned Whole Oats, Lignin Sulfonate, Dried Whey, Sodium Sulfate, Salt, Rice Bran, Sodium Sulfite, Brewers Dried Yeast, Soybean Oil, Dried Saccharomyces Cerevisiae Fermentation Product, Monocalcium Phosphate, Potassium Sulfate, Propionic Acid (Preservative), Yeast Culture, Magnesium Oxide, Sodium Bentonite, Hydrated Sodium Calcium Aluminosilicate, Aspergillus Oryzae Fermentation Extract, Potassium Chloride, Zinc Sulfate, Natural and Artificial Flavors, Magnesium Sulfate, Manganese Sulfate, Iron Oxide, Propylene Glycol, Magnesium Chloride, Extracted Streptomyces Meal, Condensed Fermentation Corn Extractives, Copper Sulfate, Vitamin A Supplement, Manganous Oxide, Kelp Meal, Vitamin D3 Supplement, Certified Color Added, Ferrous Sulfate, Ferrous Carbonate, Manganese Proteinates, Vitamin E Supplement, Zinc Proteinates, Zinc Oxide, Riboflavin Supplement, Vitamin B-12 Supplement, Calcium Pantothenate, Copper Carbonate, Copper Amino Acid Complex, Cobalt Glucoheptonate, Biotin, Mineral Oil, Choline Chloride, Corn Cob Fractions, Niacin Supplement, Potassium Sulfite, Selenium Yeast, Calcium Iodate, Pyridoxine Hydrochloride, Thiamine Mononitrate, Cobalt Carbonate, Cobalt Sulfate, Folic Acid, Cobalt Amino Acid Complex, Ethylenediamine Dihydriodide, Sodium Selenite, Ethoxyquin (Preservative), Silicon Dioxide, Citric Acid (Preservative), Tertiary Butyl Hydroquinone (Preservative).

FEEDING DIRECTIONS

SHOW CATTLE:

FOR INCREASED RATE OF WEIGHT GAIN (Growing cattle on pasture or in dry lot [stocker and feeder, and dairy and beef replacement heifers] AND Mature reproducing beef cows [on pasture in dry lot]): Hand feed or mix in total ration so as to supply 3.63 to 14.50 lbs. of this feed per head per day, thereby supplying 50 - 200 mg monensin per head daily. During first 5 days, cattle should receive no more than 100 mg monensin per day (7.25 lbs. of this feed).

CONTINUED ON BACK --->

SHOWTIME HEIFER & STEER DEVELOPER

(Back Side)

FEEDING DIRECTIONS (Continued)

FOR THE PREVENTION AND CONTROL OF COCCIDIOSIS CAUSED BY EIMERIA BOVIS AND E. ZUERNII (Cattle fed in confinement for slaughter): Feed, depending upon severity of challenge, 1.02 to 3.05 lbs of this feed per 100 lbs body weight daily, supplying 14 to 42 mg monensin per 100 lbs body weight daily. (Growing cattle on pasture or in dry lot [stocker and feeder, and dairy and beef replacement heifers] AND Mature reproducing beef cows [on pasture in dry lot]): Feed, depending upon severity of challenge, 1.02 to 3.05 lbs of this feed per 100 lbs body weight daily, supplying 14 to 42 mg monensin per 100 lbs body weight daily up to 200 mg monensin (14.50 lbs of this feed) per head daily. During first 5 days of feeding, cattle should receive no more than 100 mg monensin (7.25 lbs of this feed) per day.

(Calves[excluding veal calves]): Feed, depending upon severity of challenge, 1.02 to 7.25 lbs of this feed per 100 lbs body weight daily, supplying 14 to 100 mg monensin per 100 lbs body weight daily up to 200 mg monensin (14.50 lbs of this feed) per head daily.

FOR IMPROVED FEED EFFICIENCY (Cattle fed in confinement for slaughter): Feed continuously so as to supply 3.63 to 26.10 lbs. of this feed per head daily, thereby supplying 50 to 360 mg monensin per head daily.

CAUTION: Do not allow horses or other equines access to feed containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin-medicated cattle feed is safe for use in cattle only. Consumption by unapproved species may result in toxic reactions. Do not exceed the levels of Monensin recommended by the feeding directions, as reduced average daily gains may result. A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal.

WARNING! DO NOT FEED TO SHEEP. This feed contains added copper at levels that may be toxic to sheep.

REMINDER: Feed is perishable. Store this product in a cool, dry area away from insects. DO NOT feed moldy or insect-infested feed.

Net Weight:

50 Pounds (22.7 Kilograms)

Manufactured By :

Farmers Warehouse Company
P.O. Box 160
Keyes, California 95328
(209)632-2333

15565503.02
010-A-5820

426D