

BEST START[®]

5518 ALG HEIFER GROWER MIX(2010-15)~M

M E D I C A T E D

For Growing Dairy Cattle

WARNING! DO NOT FEED TO SHEEP. This feed contains added copper at levels that may be toxic to sheep.

CAUTION: Do not allow horses or other equines access to feed containing Monensin. **Ingestion of Monensin by equines has been fatal.** Monensin-medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. If feed refusals containing Monensin are feed to other groups of cattle, the concentration of Monensin in the refusals and amount of refusals fed could be taken into consideration to prevent Monensin overdosing.

ACTIVE DRUG INGREDIENTS

Monensin(Rumensin [®])	39.10 gm/t
Diflubenzuron	2.55 mg/lb

<i>Guaranteed Analysis</i>		<i>Ingredients</i>
Crude Protein	Min 16.00 %	Steam Flaked Corn, Soybean Meal, Steam Rolled Barley, Cane Molasses, Suncured Alfalfa Pellets Ground, Dried Plain Beet Pulp Shreds, Lactose Dried, Wheat Millrun, Gelatinized Corn Flour (Corn Starch), Ground Corn, Vegetable Oil, Glycerin, Ground Limestone, Hydrolyzed Palm Oil - Feed Grade, Saccharomyces Cerevisiae Yeast Culture, Dried Whey, Salt, Magnesium Oxide, Hydrated Sodium Calcium Aluminosilicate, Propionic Acid (Preservative), Natural and Artificial Flavors, Potassium Chloride, Ammonium Hydroxide, Corn Mill By-Product, Rice Bran, Zinc Sulfate, Mixed Tocopherols, Manganous Sulfate, Potassium Sulfate, Magnesium Sulfate, Monocalcium Phosphate, Ferrous Sulfate, Kelp Meal, Vitamin A Supplement, Vitamin D3 Supplement, Iron Oxide, Sodium Sulfate, Vitamin E Supplement, Copper Sulfate, L-Lysine Monohydrochloride, Sodium Sulfite, Corn Distillers Dried Grains With Solubles, Extracted Streptomyces Meal, Condensed Fermentation Corn Extractives, Manganous Oxide, Mineral Oil, Choline Chloride, Ferrous Carbonate, Sodium Bentonite, Lignin Sulfonate, Silicon Dioxide, Biotin Supplement, Folic Acid, Propylene Glycol, Pyridoxine Hydrochloride, Cobalt Glucoheptonate, Niacin Supplement, Calcium Pantothenate, Thiamine Mononitrate, Vitamin B-12 Supplement, Riboflavin Supplement, Calcium Iodate, Cobalt Carbonate, Sodium Selenite, Biotin.
Crude Fat	Min 2.20 %	
Crude Fiber	Max 5.20 %	
ADF	Max 7.10 %	
Calcium (Ca)	Min 0.30 %	
Calcium (Ca)	Max 0.50 %	
Phosphorous (P)	Min 0.30 %	
Sodium (Na)	Max 0.16 %	
Salt (NaCl)	Min 0.17 %	
Salt (NaCl)	Max 0.25 %	
Potassium (K)	Min 1.00 %	
Vitamin A	Min 10000 IU/lb	

CONTINUED ON BACK - - - >

BEST START[®] 5518 ALG HEIFER GROWER MIX(2010-15)~M (Back Side)

FEEDING DIRECTIONS

FOR THE PREVENTION AND CONTROL OF COCCIDIOSIS CAUSED BY EIMERIA BOVIS AND E. ZUERNII (For growing cattle): Feed, depending upon severity of challenge, 0.7 to 2.2 lbs of this feed per 100 lbs body weight daily, supplying 14 to 42 mg monensin per 100 lbs body weight daily up to 200 mg monensin (10.30 lbs of this feed) per head daily. During first 5 days of feeding, cattle should receive no more than 100 mg monensin (5.15 lbs of this feed) per day.

FOR INCREASED RATE OF WEIGHT GAIN (For growing cattle): Hand feed or mix in total ration so as to supply 2.6 to 10.30 lbs. of this feed per head per day, thereby supplying 50 - 200 mg monensin per head daily. During first 5 days, cattle should receive no more than 100 mg monensin per day (5.15 lbs. of this feed).

WARNING: A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal.

Feed this product to cattle according to your nutritionist's directions. For selenium purposes, this product should not exceed 52.63% of total ration.

REMINDER: Feed is perishable. Store this product in a cool, dry area away from insects. DO NOT feed moldy or insect-infested feed.

This feed contains an insect growth regulator, which prevents the development of house, stable, face and horn flies in the manure of treated beef and dairy cattle. Feed this product to beef or dairy cattle for continuous feeding during the fly season at a rate of 1.80 lbs per one hundred pounds of body weight per day to provide 4.54 mg Diflubenzuron per one hundred pounds of body weight per day (0.10 mg/kg of body weight/day).

Net Weight: **50 Pounds (22.7 Kilograms)**

Manufactured By :

Farmers Warehouse Company
P.O. Box 160
Keyes, California 95328
(209)632-2333

1586551829

Manufactured For:

A. L. Gilbert Company
Oakdale & Keyes, California
1-800-847-1721

832Q