



FARMERS BEST CHICK STARTER/GROWER CRUMBLES

M E D I C A T E D

As an aid in the prevention of coccidiosis

Active Drug Ingredient: Amprolium 0.0125 %

WARNING: Use as the sole source of Amprolium.

G U A R A N T E E D A N A L Y S I S

| | | | | | |
|-----------------|-----|---------|--------------|-----|--------|
| Crude Protein | Min | 20.00 % | Calcium (Ca) | Min | 0.20 % |
| Lysine | Min | 0.90 % | Calcium (Ca) | Max | 0.40 % |
| Methionine | Min | 0.30 % | Salt (NaCl) | Min | 0.30 % |
| Crude Fat | Min | 4.00 % | Salt (NaCl) | Max | 0.60 % |
| Crude Fiber | Max | 6.00 % | Sodium (Na) | Max | 0.25 % |
| Phosphorous (P) | Min | 0.40 % | | | |

I N G R E D I E N T S

Ground Corn, Soybean Meal, Safflower Meal, Calcium Carbonate, Hydrolyzed Vegetable Oils, Suncured Alfalfa Meal, Monocalcium Phosphate, Dicalcium Phosphate, Salt, DL-Methionine Hydroxy Analogue, Choline Chloride, Ferrous Sulfate, Manganous Oxide, Copper Sulfate, Ethylenediamine Dihydriodide, Zinc Oxide, Mineral Oil, Sodium Selenite, Niacin Supplement, Calcium Pantothenate, Riboflavin Supplement, Vitamin A Supplement, Vitamin E Supplement, Vitamin D3 Supplement, Vitamin B-12 Supplement, Menadione Sodium Bisulfite Complex, Ethoxyquin (Preservative), Folic Acid, Thiamine Mononitrate, Pyridoxine Hydrochloride.

F E E D I N G D I R E C T I O N S

Start the chicks eating early by sprinkling FARMERS BEST CHICK STARTER/GROWER CRUMBLES (Medicated) on paper plates or egg mapes 1/3 full of feed. Ath the end of the first week, begin feeding in chick size feed hoppers. Allow one inch of feeding space per chick. By 3rd week, double available feeding space to 2 inches per chick, and use medium size feeders. Fill feeders only 2/3 full to help prevent waste. Feed FARMERS BEST CHICK STARTER/GROWER CRUMBLES (Medicated) through the 8th week. Provide fresh, clean water at all times, allowing 1/2 inch per chick. Grit should be made available after the 2nd week. Troubles with chicks are commonly caused by overcrowding, poor sanitation, chilling, overheating or carelessness. Always buy chicks from a reliable hatchery. Brood chicks as far as possible from adult poultry to help prevent disease.

WARNING! DO NOT FEED TO SHEEP. This feed contains added copper at levels that may be toxic to sheep.

REMINDER: Feed is perishable. Store this product in a cool, dry area away from insects. DO NOT feed moldy or insect-infested feed.

Net Weight:

50 Pounds (22.7 Kilograms)

Mfd By: Farmers Warehouse Company, P.O. Box 160, Keyes, California 95328, (209)632-2333

1506962701

832Q