

**FARMERS BEST 24/18 CF
CALF MILK REPLACER
BOVA/MOS
MEDICATED**

For control of coccidiosis caused by *Eimeria bovis* and *E. zuernii* in replacement calves.

ACTIVE DRUG INGREDIENT

Lasalocid (as lasalocid sodium).....60 grams/ton

GUARANTEED ANALYSIS

Crude Protein, min.....24.00%
Crude Fat, min.....18.00%
Crude Fiber, max.....0.15%
Calcium, min.....0.75%
Calcium, max.....1.25%
Phosphorus, min.....0.70%
Vitamin A, min.....30,000 IU/LB
Vitamin D3, min.....10,000 IU/LB
Vitamin E, min.....200 IU/LB
Vitamin C, min.....225 MG/LB

INGREDIENTS

Dried Skimmed Milk, Dried Whey, Dried Whey Protein Concentrate, Dried Whey Product, Animal Fat (Preserved with BHT), Coconut Oil, Lecithin, Dried Milk Protein, Yeast Primary Dehydrated, Dicalcium Phosphate, Calcium Carbonate, Magnesium Sulfate, l-Lysine, dl-Methionine, Vitamin A Acetate, Vitamin D₃ Supplement, Vitamin E Supplement, Copper Proteinate, Manganese Proteinate, Zinc Proteinate, Iron Proteinate, Selenium Yeast, Dried Brewer's Yeast, Vitamin B₁₂ Supplement, Thiamine Mononitrate, Ascorbic Acid, Biotin, Riboflavin, d-Calcium Pantothenate, Niacinamide, Choline Chloride, Menadione Sodium Bisulfite Complex (Source of Vitamin K₃), Folic Acid, Pyridoxine Hydrochloride, Calcium Iodate, Zinc Sulfate, Ferrous Sulfate, Manganese Sulfate, Copper Sulfate, Cobalt Sulfate, Sodium Silico Aluminate, Polyoxyethylene Glycol (400) Mono and Dioleates, Artificial Flavor.

WARNING: A WITHDRAWAL PERIOD HAS NOT BEEN ESTABLISHED FOR LASALOCID IN PRE-RUMINATING CALVES. DO NOT USE IN CALVES TO BE PROCESSED FOR VEAL.

CAUTION: DO NOT ALLOW HORSES OR OTHER EQUINES ACCESS TO FEEDS CONTAINING LASALOCID AS INGESTION MAY BE FATAL. THE SAFETY OF LASALOCID IN UNAPPROVED SPECIES HAS NOT BEEN ESTABLISHED.

See Feeding Instructions on Reverse Side

This feed was produced in a facility certified in the American Feed Industry Association's Safe Feed/Safe Food Certification Program; for details go to:

www.safefeedsafefood.org.



FEEDING INSTRUCTIONS

Hand feed at a rate of 1 mg of lasalocid per 2.2 pounds body weight per day. Thoroughly mix this milk replacer powder with warm water and feed immediately. When preparing large batches (i.e. anything other than individual calf servings) keep the mixture agitated while dispensing into individual feeding containers to prevent settling. Each pound of this milk replacer powder contains 30 mg of lasalocid; one pound of this milk replacer powder will medicate 66 pounds of body weight.

It is recommended that calves be given 2 to 3 quarts of good quality colostrum as soon as possible after birth, and another 2 to 3 quarts within the first 24 hours of life. Second milking colostrum may be fed afterwards for up to 3 days. Starting on the 4th day, feed a high quality milk replacer according to the following directions.

MIXING

Into 3 quarts of warm (120°F) clean water, add 12 ounces (1 cup)* of milk replacer powder and mix thoroughly. For larger batches, mix 16 ounces of milk replacer powder per gallon of warm water.

FEEDING

Feed the reconstituted milk replacer at a rate of 3 quarts per 100 pounds of body weight twice daily (6 quarts per 100 pounds of bodyweight per day). Large breed calves generally require 3 to 3-1/2 quarts per feeding twice daily, while smaller breed calves may need 2-1/2 to 3 quarts twice daily.

MANAGEMENT SUGGESTIONS

- For best mixing results, sprinkle milk replacer powder onto water.
- Use of a wire whisk or mechanical agitation provides good mixing.
- Feed reconstituted milk replacer shortly after mixing.
- Feeding and mixing equipment should be cleaned and sanitized after each use.
- Clean, fresh water should be available at all times.
- Calves should have access to a quality texturized feed after 4 days of age.
- Hay should not be offered until after calves are weaned.
- Wean calves when they are consuming 1-1/2 pounds of starter feed per day for 3 consecutive days.

* The measuring cup provided in the bag holds approximately 12 ounces of milk replacer powder. For greater accuracy, a suitable scale is recommended.